

Exhibit B

[VIEW IN FRAME](#)

Pharmaceutical Abbreviations

This web page contains abbreviations and symbols that are commonly used by doctors when writing a prescription. In addition to the standard pharmaceutical abbreviations, we've included a few abbreviations used with clinical trials.

If you don't find what you are looking for here, try the [AEGIS Medical Glossary](#).

| | |
|----------------|---|
| a.c. | before eating (ante cibum) |
| ad lib. | as much as wanted (ad libitum) |
| b.i.d. | two times a day (bis in die) |
| b.i.w. | two times a week |
| DOT | direct observational therapy |
| d.t.d.no. ____ | give ____ doses (dentur talis doses numus ____) |
| e.m.p. | in the manner prescribed (ex modo praescripto) |
| ft. | make (fac, fiat, fiant) |
| gm. | gram |
| gtt. | drop, drops (gutta, guttae) |
| h. | hour |
| h.s. | at bedtime (hora somni) |
| ID | intradermal |
| IL | intralesional |
| IM | intramuscular |
| IN | intranasal |
| IND | investigational new drug (US FDA designation) |
| IU | international unit |
| IV | intravenous |
| kg. | kilogram |
| m. | month |
| m.ft. | mix and make (miscre et fiat) |
| µg | microgram |
| mg. | milligram |
| ml. | milliliter |
| m.t. | send like this (mitte talis) |
| MTD | |

| | |
|-----------------|--|
| | maximum tolerated dose |
| n.r. (non rep.) | do not repeat (non repetatur) |
| o.d. | right eye (oculo dextro) |
| o.s. | left eye (oculo sinistro) |
| o.u. | each eye (oculo utro) |
| p. | after (post) |
| p.c. | after meals (post cibum) |
| pcg. | picogram |
| p.o. | taken by mouth |
| p.r.n. | as necessary (pro re nata) |
| q. ___ h. | every ___ hours (quaque 4 horae) ("q4h" would mean "every four hours") |
| q.d. | every day (quaque die) |
| q. ___ d. | every ___ days ("q9d" would mean "every nine days") |
| q.i.d. | four times a day (quarter in die) |
| q.i.w. | four times a week |
| q.m. | every month |
| q.o.d. | every other day |
| q.o.w. | every other week |
| q.s. ad | a sufficient quantity to make (quantum sufficiat ad) |
| q.w. | every week |
| SC | subcutaneous |
| s.o.s. | if needed (si opus sit) |
| ss | one half (semis) |
| stat. | immediately (statim) |
| t.i.d. | three times a day (ter in die) |
| t.i.w. | three times a week |
| tal. | of such (talis) |
| ut dict. | as directed (ut dictum) |
| WNL | within normal limits |

This information is designed to support, not replace, the relationship that exists between you and your doctor.
©1998. AEGIS.